Express Mail No.: ET437641649 US

Sheet 1 of 3

Form PTO-1449 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE				ATTY. DOCKET NO. 2804-J		serial noto be assigned—		
FATERI AND TRADEWARK OFFICE								
INFORMATION DISCLOSURE CITATION			APPLICANT		Divisional of 09/628,126			
			IATION	Raymond G. Goodwin et al.,				
				FILING DATE		GROUP		
(Use several sheets if	necessary)	October 2, 299	93	to be a	ssigne	d
			U.S. PATENT	DOCUMENTS	,			
EXAMINER	DOCUMENT NUMBER	DATE		NAME	CLASS			G DATE
SJH	5,165,923	11/24/92	Thorpe et a	orpe et al.				
								
1		FO	REIGN PATE	NT DOCUMENTS	1	_l	L <u></u>	
	DOCUMENT NUMBER	DATE		COUNTRY		SUBCLASS	TRANS YES	LATION NO
SJH	WO 93/10232	5/27/93	PCT				x	
SJH	WO 91/07437	5/30/91	PCT					x
SJH	WO 87/07144	12/3/87	PCT					
SJH	EP 0 412050 A1	2/6/91	EP					
	OTHER DOC	uments (Including Auth	or, Title, Date, Pertinen	t Pages. E	itc.)	.,	
SJH	OTMER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.) Armitage et al., "Molecular and Biological Characterization of a Murine Ligand for CD40," Nature 357: 80-82, 1992.							
S.TH	Zsebo et al., "Stem Cell Factor is Encoded at the SI Locus of the Mouse and is the Ligand for the c-kit Tyrosine Kinase Receptor," Cell 63: 213-224, 1990.							
	Schwarting et al., "Cluster report: CD30," Leucocyte Typing IV: White Cell Differentiation Antigens: p 419-422, Oxford University Press, 1989.							
	Berenbeck et al., "Detection of soluble Hodgkin-associated CD30 antigen in the sera of patients with Hodgkin's lymphoma," Leucocyte Typing IV: White Cell Differentiation						of	
							!	
	Antigens: p 425, Oxford University Press, 1989. Dallenbach et al., "Soluble CD30 antigen in the sera of patients with adult T-cell							
	lymphoma/leukaemia (ATL): a marker for disease activity," Leucocyte Typing IV: White						ite	
	Cell Differentiation Antigens: p. 426-428, Oxford University Press, 1989.							
	Smith et al., "CD30 Antigen, a Marker for Hodgkin's Lymphoma, Is a Receptor Whose Ligand Defines an Emerging Family of Cytokines with Homology to TNF," Cell 73: 1349-							
	1360, 1993.	30 Ligand	: Molecular	Cloning and Pathobi	inlogical	Role in Cl	D30-	
	Gruss et al., "CD30 Ligand: Molecular Cloning and Pathobiological Role in CD30-Positive Malignant Lymphomas," presented at Keystone Symposium on B and T Cell							
	Lymphomas, Copper Mountain, CO, April 17-24, 1993.							
	Loh et al., "Polymerase Chain Reaction with Single-Sided Specificity: Analysis of T Cell							
	Receptor δ Chain," Science 243: 217-220, 1989.							
SJH	Falini et al., "Response of refractory Hodgkin's disease to monoclonal anti-CD30 immunotoxin," Lancet 339: 1195-1196, 1992.							
	inimianotoxin, 2	Milet 337		DATE CONSIDERED				
EXAMINER	/Sheela J. Huff/				08	/17/2006		
*EXAMINER: I not in conform	nitial if citation considered ance and not considered	d, whether or . Include co	r not citation is i	in conformance with MP with next communication	EP 609; D	raw line throu	igh citati	ion if

<u>:</u>				· · · · · · · · · · · · · · · · · · ·					
F	rm PTO-	·1449	U.S. DEPARTMENT OF COMMERCE	ATTY, DOCKET NO.	SERIAL NO.				
	PATENT AND TRADEMARK OFFICE		2804-J	to be assigned					
					Divisional of 09/628,126				
	INFORM	MATION [DISCLOSURE CITATION	APPLICANT					
	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			Raymond G. Goodwin, et al.					
				FILING DATE	GROUP				
L		(Suppl	lemental Sheet)	October 2, 2003	to be assigned				
		0.	THER DOCUMENTS (Including Au	thor, Title, Date, Pertinent Page:	s, Etc.)				
		da Costa et al., "Immunoscintigraphy in Hodgkin's disease and anaplastic large cell							
	SJH		lymphomas: Results in 18 patients using the iodine radiolabeled monoclonal antibody HRS-						
L			Onc. 3 (Suppl. 4): \$53-\$57, 199	7, 1992.					
		Engert e	ert et al., "Evaluation of Ricin A Chain-containing Immunotoxins Directed against the						
		CD30 Antigen as Potential Reagents for the Treatment of Hodgkin's Disease," Cancer Res.							
L		<i>50:</i> 84-8	88, 1990.						
			et al., "Experimental therapy in H	Hodgkin's disease," Ann. Or	nc. 3(Supp.4): S97-S100,				
$ldsymbol{ld}}}}}}$		1992.							
			al., "Three new lymphoid activa		yping III: White Cell				
<u>_</u>			ttiation Antigens, p. 574, Oxford						
	ł l		ting et al., "BER-H2: a new mon						
			kin's disease in formaldehyde-fi						
<u> </u>			ferentiation Antigens, pp. 574-5						
			n et al., "The Role of Ki-1 Antig						
			ernberg Cells, Ki-1 Lymphoma (exytes," Blood 80(10)				
<u> </u>			305a, abstract no. 1211, Nov. 15						
			l., "EBV and Tumor Progression		Cell Lymphoma (ALCL),"				
			0(10) Supp. 1: 445a, Abstract no						
			n et al., "The TNF/NGF Superfa						
			n., Keystone Symposia on Molec	ular & Cellular Biology, Su	appl. 17B, abstract F020,				
	_		Feb. 10, 1993.						
			and Hamilton-Dutoit, "Ki-1 (Cl						
<u>_</u>			Embryonal Carcinoma," Am. J.						
			ing et al., "BER-H2: A New Ant		Antibody Directed at a				
$oxed{oxed}$	┦—	Formol-Resistant Epitope," Blood 74: 1678-1689, 1989.							
		Andreesen et al., "Human Macrophages Can Express the Hodgkin's Cell- Associated Antigen							
<u>_</u>			D30)," Am. J. Path. 134: 187-19						
			rsheimer and Möller, "Expression	n of Ki-1 Antigen (CD30) i	n Mesenchymal Tumors,"				
		Cancer 66(8): 1732-1737, 1990.							
		Gianotti et al., "Primary Cutaneous Pleomorphic T-Cell Lymphoma Expressing CD30							
<u> </u>	╀	Antigen," Am. J. Dermatopath. 13(5): 503-508, 1991.							
		Burns and Dardick, "Ki-1 Positive Non-Hodgkin's Lymphomas," Am. J. Clin. Path. 93(3):							
<u> </u>		327-332, 1990.							
<u> </u>		Piris et al., "CD30 expression in non-Hodgkin's lymphoma," Histopath. 17: 211-218, 1990.							
		Miettinen, "CD30 Distribution," Arch. Pathol. Lab. Med. 116: 1197-1201, 1992.							
`	<u> </u>		al., "CD30 expression in follicula						
Maeda and Takahashi, "Characterization of skin infiltrating cells i					adult T-cell				
leukaemia/lymphoma (ATLL): clinical, histological a					tochemical studies on				
eight cases," Brit J. Derm. 121: 603-612, 1989.									
EX	AMINER	(-1 1 -		DATE CONSIDERED	20/17/2006				
/Sheela J. Huff/ 08/17/2006									
*EX	CAMINER: Ir	nitial if citatio	on considered, whether or not citation is	s in conformance with MPEP 609	: Draw line through citation if				
no	t in conform	ance and no	ot considered. Include copy of this form	n with next communication to apr	olicant.				

·							
Form PTC	1449	U.S. DEPARTMENT OF COMMERCE	ATTY, DOCKET NO.	SERIAL NO.			
		PATENT AND TRADEMARK OFFICE	2804-J	to be assigned			
INFORMATION DISCLOSURE CITATION				Divisional of 09/628,126			
				APPLICANT			
			Raymond G. Goodwin, et al.				
			FILING DATE	GROUP			
	(Suppl	lemental Sheet)	October 2, 2003	To be assigned			
ļ		THER DOCUMENTS (Including Au					
SJH	Positive	t al., "Follicular Lymphoid Hyp Lymphocytes," Am. J. Dermato	opath. 11(4): 345-352, 1989	·			
	Aruffo and Seed, "Molecular Cloning of a CD28 cDNA by a High Efficiency COS Cell Expression System," <i>Proc. Nat. Acad. Sci. USA 84</i> : 8573-8577, 1987.						
	Flanagan and Leder, "The kit Ligand: A Cell Surface Molecule Altered in Steel Mutant Fibroblasts," Cell 63: 185-194, 1990.						
	Josimovic-Alasevic et al., "Ki-1 (CD30) Antigen is Released by Ki-1 Positive Tumor Cells in						
	vitro and in vivo," Eur. J. Immunol. 19: 157-162, 1989. Froese et al., "Biochemical Characterization and Biosynthesis of the Ki-1 Antigen in Hodgkin-Derived and Virus-Transformed Human B and T Lymphoid Cell Lines," J.						
	Immunol. 139(6): 2081-2087, 1987.						
	Shohat et al., "Inhibition of Cell Growth Mediated by Plasmids Encoding p53 Antisense," Oncogene 1: 277-283, 1987.						
	Riordan and Martin, "Oligonucleotide-based Therapeutics," <i>Nature 350:</i> 442-443, 1991.						
	Pfreundschuh et al., "In-vivo-Imaging von Hodgkin-Lymphomen mit monoklonalen						
	Antikörpern," Onkologie 12: Supp. 1, 30-33, 1989.						
	Carde et al., "Immunoscintigraphy of Hodgkin's Disease: In Vivo use of Radiolabelled						
	Monoclonal Antibodies Derived from Hodgkin Cell Lines," Eur. J. Cancer 26(4): 474-479, 1990.						
	Stein et al., "The Expression of the Hodgkin's Disease Associated Antigen Ki-1 in Reactive						
		plastic Lymphoid Tissue: Evide					
 	Malignancies Are Derived from Activated Lymphoid Cells," Blood 66(4): 848-853, 1985.						
	Andreesen, et al., "A Hodgkin Cell-Specific Antigen is Expressed on a Subset of Auto-and Alloactivated T (Helper) Lymphoblasts," <i>Blood 63(6):</i> 1299-1302, 1984. Klein, "Epstein-Barr Virus-Carrying Cells in Hodgkin's Disease," <i>Blood 80(2):</i> 299-301, 1992.						
	Poiesz, et al., "Detection and isolation of type C retrovirus particles from fresh and cultured						
	lumphocytes of a patient with cutaneous T-cell lymphoma," <i>Proc. Natl. Acad. Sci. USA</i> 77(12): 7415-7419, 1980.						
	Schwab et al., "Production of a monoclonal antibody specific for Hodgkin and Sternberg-						
	Reed cells of Hodgkin's disease and a subset of normal lymphoid cells," <i>Nature 299:</i> 65-67, 1982.						
	Stein et al., "Identification of Hodgkin and Sternberg-Reed Cells as a Unique Cell Type Derived from a Newly Detected Small-Cell Population," Intl. J. Cancer 30: 445-459, 1982.						
	Durkop et al., "Molecular Cloning and Expression of a New Member of the Nerve Growth						
	Factor Receptor Family that is Characteristic for Hodgkin's Disease," Cell 68: 421-427, 1992.						
SJH	Capon et al. "Designing CD4 Immunoadhesins for AIDS Therapy" Nature 337: 525-531						
EXAMINER	/Cheal=	J. Huff/	DATE CONSIDERED	8/17/2006			
*EVANUESE							
not in confor	*EXAMINER: Initial if citation considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.						